



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

lowing periodic phases were noted, as forming an annual cycle: (1) A winter phase with free diatoms in abundance; (2) a spring phase with *Spirogyra* dominant; (3) a summer phase with *Cladophora* dominant and epiphytes abundant; and (4) an autumn phase in which many of the earlier forms become active again after a latent period. These include *Spirogyra*, *Edogonium*, or other forms. The authors classify and discuss the factors that operate to produce these changes as seasonal, irregular, and correlated. This kind of continued work with small bodies of water is needed in this country; and our own members can make valuable contributions to the ecology of our micro-organisms by such investigations in the neighborhood of their homes.

DISTRIBUTION OF ROTIFERS

In the *Journal* of the *Queckett Microscopical Club*, C. F. Rousselet offers a discussion of the distribution of Rotifers. He calls attention to the fact that most species have quite a cosmopolitan range; and that no continent or climatic zone can really claim a peculiar rotatorian fauna. Even the rarer species are reported from widely separated regions of the earth. These facts are of course to be coupled with the ease with which Rotifers are transported. Some of them, as is known, are even capable of being dried out and of resuming life when conditions become favorable. Even in species in which this is not true there are resting eggs that resist both cold and drouth. The fact that the habitat of many Rotifers is such that they or their eggs are liable to be dried up once or oftener each year, and thus to be committed to the winds, is likewise an important factor.

RELATION OF VITALITY OF PARAMECIUM TO CONSTANCY OF SURROUNDINGS

Mr. L. L. Woodruff states, in the *Biological Bulletin*, that Paramecia kept for generations in a reasonably constant culture medium undergo cyclical changes in protoplasmic vitality; and finally die from internal causes. If, however, the culture medium is kept changing in such a way as to disturb this cycle, the lowering of vitality may be prevented and the protoplasmic life may be continued without any apparent decrease of vitality. Possibly the cycle leading to senility may even be eliminated altogether in this way.